784.8669

RESOURCE GUIDE



MICHIGAN TOBACCO QUITLINE 1.800 **QUIT.NOW**


Get FREE Confidential **Counseling & Support**

DOUBLE your chances of quitting.

> Call Now **1.800.QUIT.NOW**

www Or Enroll Online michigan.quitlogix.org





TREATMENT OPTIONSQUICK GUIDE



PATCH

21 mg If you smoke more than 10 cigarettes per day

STEP1

Select your starting dose.*



STEP 2

STEP 3

STEP 4

Weeks 1-6

Place the patch on a clean, dry, hairless spot on your upper body or

Reduce dosage over time.*

7 mg

outer arm. Press for 10 seconds.

Replace patch every 16-24 hours.

STEP 3

options!

GUM

STEP 1

STEP 2

Choose a flavor.

There are many

Select your strength.*

Chew slowly until the taste becomes strong or tingles, place the gum between your cheek and gums for 1 minute or until tingle is gone. Repeat this chew and park method for 30 minutes.



STEP 4

2 mg

Use the gum whenever you have the urge to smoke as part of a quit immediately or quit gradually plan.*

PRESCRIPTION ONLY

This option

helps keep

your hands busy!

If you smoke

more than 25

cigarettes per day

If you smoke 25

per day

or less cigarettes

LOZENGE

STEP 3

1-2

hours

Weeks

1-6

STEP1
Select your strength*

Select your strength.*

STEP 2

Reduce usage over time.*

2-4

hours

Weeks

7-9

Use 1 lozenge every...

Slowly dissolve the lozenge in your mouth

(between cheek and gums), while occasionally

moving it from one side of your mouth to the

other. The lozenge will take approximately 10-20

minutes to fully dissolve. Do not chew or swallow.

4 mg If you smoke within 30 mins of waking up



smoke NASAL SPRAY

STEP 1

Blow your nose to clear it. Prime pump with 6-8 sprays prior to first use.



7 mg

Weeks 9-10

PRESCRIPTION ONLY

1 dose = 2 sprays (one in each nostril)

STEP 2

Tilt your head back and spray once in each nostril while breathing through your mouth. Avoid sniffing or inhaling. Wait 2-3 minutes before blowing your nose.

14 mg

Weeks 7-8

STEP 3

Reduce dosage over time.*



INHALER

STEP 1

Insert a cartridge into inhaler.

Push and twist mouthpiece back on.

STEP 2

Puff in short breaths through the mouthpiece like you are sipping from a straw. Cartridges last for about **20 minutes** of active use, but this can be broken up into multiple shorter uses.

STEP 3

When the cartridge is empty, remove from the inhaler and discard.

STEP 4

Reduce usage over time.*

6-16 cartridges per day

Reduce gradually

Weeks 1-12

Weeks 13-24

24

hours

Weeks

13-14

4-8

hours

Weeks

10-12

^{*} Consult your doctor to determine the appropriate treatment options and dosing.